

REMARKS ON D-DIMENSIONAL TSP OPTIMAL TOUR LENGTH BEHAVIOUR

ANDREI G. YANEF

ABSTRACT. The well-known $O(n^{1-1/d})$ behaviour of the optimal tour length for TSP in d-dimensional Cartesian space causes breaches of the triangle inequality. Other practical inadequacies of this model are discussed, including its use as basis for approximation of the TSP optimal tour length or bounds derivations, which I attempt to remedy.

1. INTRODUCTION

Studies of the behaviour of the optimal tour value in respect of the TSP has been in place since the fifties. Considerate work has been done with the $O(n^{1-1/d})$ model of behaviour based on results of Beardwood, Halton and Hammersley [1]. The approach taken here is one of simple real analysis and some of the intermediate results correspond with those of Snyder and Steele [8] and Bern and Eppstein [2].

As pointed out by Bern and Eppstein [2], an a priori bound on a geometric graph is a bound that depends on the assumption that all vertices lie within a given container. Rather than bounds, I consider the behaviour of the optimal tour length of a geometric graph in a 2D rectangular region, which is transformed to a square and then simplified (extended) to d-dimensional cube. This is compared to the behaviour of the MST length in the same container space.

The problem of studying the behaviour or approximation of the optimal tour length of TSP instances ($\|O\|$) can be illustrated by an example: Given two TSP instances with substantially different number of vertices have the same optimal tour length.

It is shown that the general behaviour of $\|O\|$ does not depend on the number of vertices. I also give experimental evidence in support of this using a small set of TSPLIB [5] instances, and conclude that the commonly cited $O(n^{1-1/d})$ behaviour is inadequate in both description of behaviour (nor useful bounds) of $\|O\|$. A simple alternative is suggested.

The present is an extended version of a mention in [9].

2. TERMS AND DEFINITIONS

The Minimum TSP in Cartesian space is a special case of the Minimum Weight Hamiltonian Cycle problem. Both are specified as follows:

Date: August 14, 2002 and, in revised form, September 18, 2002.
1991 Mathematics Subject Classification. ACM F.2.2 ; G.2.2.
Key words and phrases. TSP, MST, optimal tour length, bounds.
 E.T.C.2/2002.

P# 1 (Minimum Hamiltonian Cycle). *Given $G = (V, E)$ with a cost function $d : E \rightarrow R$, find solution $O = (V_O, E_O) \subseteq G$ such that*

- (a) *O is a Hamiltonian cycle;*
- (b) *$\|O\| = \sum_{e \in E_O} d(e)$ is minimum;*

P# 2 (Minimum TSP in Cartesian Space). *P#1 and*

- (a) *$\forall v : V \cdot \text{coords}(v) \in R^d$ and*
- (b) *$\forall (u, v) : E \cdot d(u, v) = |\text{coords}(u) - \text{coords}(v)|$*

where $\text{coords} : V \rightarrow R^d$ are Cartesian.

I (unnecessarilly) split the problem specification this way for my personal convenience in what follows (separating construction and geometric aspects of the problems), and for the following observations (from P#1):

- (a) $|V| = |V_O| = |E_O|$;
- (b) Solution bounds:

$$(1) \quad nw_0 \leq \|O\| \leq nw_1$$

$$\begin{aligned} \text{where } n &= |E_O|, \\ w_0 &= \min\{d(e) | e \in E\} \wedge w_1 = \max\{d(e) | e \in E\} \end{aligned}$$

The notation used is fairly standard. I have attempted to maintain consistent naming, e.g. $Oabc_i$ is the value (approximate) of the exact $\|O\|$, and the subscript i is the number of 'alternative' formulation of $Oabc$, e.g. Ob_1 is an alternative to Ob_2 .

3. TWO SIMILAR FORMULATIONS OF D-DIMENSIONAL $\|O\|$ BEHAVIOUR

Consider a rectangle (being a container space of the TSP graph and respective tour) and that we have a uniform grid tight within (and on) that rectangle with size a by b points, assuming without loss that a is even. The optimal tour value and the diagonal of the rectangle are:

$$(2) \quad \|O\| = a \cdot b \cdot w_0 = n \cdot w_0$$

$$(3) \quad w_1 = w_0 \sqrt{(a-1)^2 + (b-1)^2}$$

Dividing:

$$(4) \quad \frac{\|O\|}{w_1} = \frac{a \cdot b}{\sqrt{(a-1)^2 + (b-1)^2}}$$

If $a = b$ than we have a square, and $\|O\|/w_1 = n / ((\sqrt{n} - 1) \sqrt{2})$. To carry this to Cartesian d-dimensions, (2) and (3) are changed to $O_1 = w_0 \prod_{i=1}^d a_i = n \cdot w_0$ and $w_1 = w_0 \left(\sum_{i=1}^d (a_i - 1)^2 \right)^{1/2}$. Assuming $\forall i, j : 1 \dots d \cdot a_i = a_j$ then

$$\frac{O_1}{w_1} = \frac{n}{(n^{1/d} - 1) \sqrt{d}}$$

Note that O_1 differs from the commonly cited $O(n^{1-1/d})$ optimal tour length behaviour, which can be formulated by taking $w_1 = w_0 \left(\sum_{i=1 \dots d} a_i^2 \right)^{1/2}$ (under the assumption that a is large), then:

$$\frac{O_2}{w_1} = \frac{n^{1-1/d}}{\sqrt{d}}$$

It should be noted that in the unit d-dimensional cube $w_1 = \sqrt{d}$.

For a metric (non-topological) graph $\|O\| \geq 2w_1$ must hold [6]. Both O_1 and O_2 fail to preserve the triangle inequality when $d = 1$, due to the "spatial" nature of the derivations (no container). It is also evident that O_1 preserves the triangle inequality for any $n \geq 2$, whereas O_2 fails in doing so when $n^{1-1/d} \leq 2\sqrt{d}$. (There is nothing to prevent such instances from occurring.) The ratio of the two estimates as $d \rightarrow \infty$ is:

$$(5) \quad e(n, d) = \frac{O_2}{O_1} = 1 - \frac{1}{n^{1/d}} \rightarrow 0$$

The divergent behaviour of O_1 and O_2 is fixed by imposing the condition $n \geq 2^d + k$, $k \geq 0$ in order to have a valid d-dimensional cube, and $1/2 \leq e(n, d) \leq 1$. Either O_1 or O_2 will fail in the description (if they are assumed to be descriptive) of the optimal tour length behaviour (see Fig.1).

The MST length behaviour ($\|T\|$) is analogous [7] to that of the TSP in a grid: $\|T\| = (n-1) \cdot w_0$ and $T_1/w_1 = (n-1) / ((n^{1/d} - 1)\sqrt{d})$. $\|O\|$ can be expressed in terms of $\|T\|$ (treated as known value computed exactly by Prim's algorithm, say):

$$\begin{aligned} Ot_1 &= \|T\| + \frac{w_1}{(n^{1/d} - 1)\sqrt{d}} \\ Ot_2 &= \|T\| + \frac{w_1}{n^{1/d}\sqrt{d}} \end{aligned}$$

As a matter of fact:

$$(6) \quad \|O\| \geq \|T\| + w_0$$

$$(7) \quad (2\|T\| \geq) \|O\| \geq \|T\| \frac{n}{n-1}$$

It is the case that for any metric graph $\|T\|$ accounts for at least 50% of $\|O\|$ and can be considered a major term in any attempted approximation. It is obvious that the second term in Ot_1 may well lead to $w_0 > w_1 / ((n^{1/d} - 1)\sqrt{d})$, as $n \rightarrow \infty$, and ultimately $Ot_1 \rightarrow \|T\|$ (so will Ot_2), which cannot be. This suggests that $\|O\|$ does not depend on the number of vertices. Moreover, the $e(n, d)$ term also remains in the relative difference in estimates for the remaining 50% of $\|O\|$, as seen from Ot_1 and Ot_2 . In other words, if $\|O\|$ were to depend on the number of vertices the following must be true:

$$e(n, d) = \frac{Ot_2 - \|T\|}{Ot_1 - \|T\|} = \frac{O_2}{O_1} \wedge \|T\| \neq 0$$

Obviously this is absurd, unless $Ot_1 = Ot_2 \wedge O_1 = O_2$, which is not the case. Thus, it is difficult to contemplate using O_1 or O_2 (see Fig.2) as basis for behaviour description of $\|O\|$. The Ot_i 's are slightly better (see Fig.3), exhibiting an almost "bounding" behaviour, for which we already have (6) and (7) and the HK bound (see Fig.5).

From the above one can conclude that n can do more harm than good in the analysis of $\|O\|$. Setting $n = 2^d$ (see above), yet another approximation (see Fig.4 and Table.1) to $\|O\|$ is obtained:

$$Otc_1 = \|T\| + \frac{w_1}{\sqrt{d}} \left(= Otc_2 + \frac{w_1}{2\sqrt{d}} \right)$$

X	$\min \epsilon$	$\max \epsilon$	$\max \epsilon - \min \epsilon$	ϵ_{rms}
O_1	0	1668.14	1668.14	308.96
O_2	-39.69	1632.05	1671.74	299.91
Ot_1	-49.26	0.76	50.02	14.16
Ot_2	-49.28	0.75	50.02	15.00
Otc_1	-14.64	35.27	49.91	7.75
Otc_2	-32.32	17.61	49.93	11.14
HK^*	-9.73	0.55	10.28	2.18
$\ T\ + w_0$	-49.98	0	49.98	16.25
$\ T\ \frac{n}{n-1}$	-49.98	0	49.98	15.48

TABLE 1. Relative error ranges for the various formulas. $\epsilon = 100 \cdot (X/\|O\| - 1) \%$. (The largest HK^* value +0.55 is possibly due to a misprint in [3])

3.1. Test Data. The test data consist of 85 instances in total of sizes $4 \leq n \leq 2560$, I have used 17 instances of my own and 68 instances from TSPLIB [5] of type EUC_2D, CEIL and ATT. The three norms were treated as EUC_2D without rounding. ATT are treated in the same way, as tour values were multiplied by $\sqrt{10}$ to bring them in line with unrounded EUC_2D. The HK-bound values are taken from [3]. There were 5 TSPLIB with published shortest tour, for which I have been able to find shorter tours. These are given in Table.3. The HK bound is greater than the actual tour length for *tsp225* and *att48* by 0.55% and 0.01% (see Table.3), respectively, possibly due to a typographical error (or the use of initial rounding in computing the HK relaxation). I do not consider accuracy (see Table.2) in this analysis of major importance, as for the 50 instances for which the optimal tour itself is not published, the data can be fixed (by point displacement) so the reported values are accurate (This is not the case with HK bound).

<i>Instances</i>	<i>Size</i>	<i>O</i>	$\ O\ $	<i>HK</i>
17	$4 \leq n \leq 2560$	<i>known</i>	<i>exact</i>	–
18	$48 \leq n \leq 2392$	<i>published</i>	<i>exact</i>	<i>reported</i>
50	$99 \leq n \leq 2319$	<i>reported</i>	<i>reported</i>	<i>reported</i>

TABLE 2. TSP instances

4. CONCLUSIONS

Inadequacies of $O(n^{1-1/d})$ and similar formulations are demonstrated. A probabilistic or statistical [4] argument may show otherwise.

<i>Instances</i>	$\ O_{reported}\ $	<i>HK</i>	$\ O_{published}\ $	$\ O_{shorter}\ $
<i>eil51</i>	426	422.4	429.983307	429.117879
<i>st70</i>	675	670.9	678.597412	677.194473
<i>eil101</i>	629	627.3	642.309509	640.42177
<i>ch130</i>	6110	6074.6	6110.859375	6110.722217
<i>ch150</i>	6528	6486.6	6532.282227	6530.906184
<i>att48</i>	10628	10602.1	10601.1282	=
<i>tsp225</i>	3919	3880.3	3858.999064	=

TABLE 3. Shorter tours. Values are as reported in [3] and computed from published tours in [5]

The optimal tour length does not depend on the number of vertices - consider again a grid example in a square: Let's suppose that we have found an optimal tour $S = (V_S, E_S)$ from $G = (V, E)$ (a comb-like shape, say) and insert (in G) exactly one zillion equidistant points along each arc in $E_S \subseteq E$ of the optimal tour S to get G_Z . If S_Z is an optimal tour of G_Z then $\|S\| = \|S_Z\|$. Such point insertion is not inconsequential - the number of optimal solutions may be reduced, i.e. G_Z has a lesser number of optimal solutions than G .

I make no claims as to quality (if used as estimates) of the above given formulae, as I would not use any of them (see Table.1) as an estimate of $\|O\|$. O_1 is an upper (easily proven so) bound, for any $n \geq 2$, however running the Nearest town algorithm [6] to get us an upper bound within a factor of two is simpler. A simple derandomisation of the same algorithm yields an approximate solution within a factor of $\sqrt{2}$ (provable under a geometric argument) for P#2 in the plane.

REFERENCES

1. J. Beardwood, J. H. Halton, and J. M. Hammersley, *The shortest path through many points*, Proceedings of Cambridge Philosophical Society, vol. 55, 1959, pp. 299–327.
2. Marshall Bern and David Eppstein, *Worst-case bounds for subadditive geometric graphs*, Proceedings of the ninth annual symposium on Computational geometry, ACM Press, 1993, pp. 183–188.
3. Keld Helsgaun, *An effective implementation of the lin-kernighan traveling salesman heuristic*, Writings on Computer Science 81/1998, Roskilde University, Computer Science Department, Fanstord, California, October 1998, <http://www.dat.ruc.dk/~keld/research/LKH/>.
4. D. S. Johnson, L. A. McGeoch, and E. E. Rothberg, *Asymptotic experimental analysis for the held-karp traveling salesman bound*, Proceedings of the seventh annual ACM-SIAM symposium on Discrete algorithms, ACM Press, 1996, pp. 341–350.
5. Gerhardt Reinelt, *Tsplib'95*, Tech. report, Institute fur Informatik, University of Heidelberg, December 2001, <http://www.iwr.uni-heidelberg.de/groups/comopt/software/TSPLIB95/>.
6. Edward M. Reingold, Jurg Nievergelt, and Narsingh Deo, *Combinatorial algorithms - theory and practice*, russian translation ed., Prentice-Hall and MIR, NJ, Moscow, 1980.
7. Timothy Law Snyder, *On minimal rectilinear steiner trees in all dimensions*, Proceedings of the sixth annual symposium on Computational geometry, ACM Press, 1990, pp. 311–320.
8. Timothy Law Snyder and J. Michael Steele, *A priori inequalities for the euclidean traveling salesman*, Proceedings of the eighth annual symposium on Computational geometry, ACM Press, 1992, pp. 344–349.




FIGURE 1. Plots of $e(2^d + k, d)$, for $k = \log_2 d$, $k = d^{\log_2 d}$, $k = d^d$, and $k = d^{d+1}$, for $d = 1 \dots 100$.




FIGURE 2. Point histograms of relative error of O_1 and O_2 - 85 instances

9. Andrei Yaneff, *On the approximation of the optimal tour length for the travelling salesman's problem*, Talk given at the ACM-SIAM Conference on Discrete Mathematics, August 2002.

E.T.CONSULTING, 6 BUNTOVNIK STREET, SOFIA, 1421, BULGARIA
E-mail address: agy@techno-link.com




FIGURE 3. Point histograms of relative error of Ot_1 , Ot_2 and HK bound (-line used for clarity)




FIGURE 4. Point histograms of relative error of Otc_1 and Otc_2




FIGURE 5. Point histograms of the lower bounds $\|T\|+w_0$, $\|T\|\frac{n}{n-1}$ and HK bound (-line used for clarity)